



HAL
open science

Dispelling myths on external factors affecting post-thawing motility of canine spermatozoa: a retrospective study on 1242 samples

Alessandra Paganotto, Camille Langlade, Emilie Rosset, S. Buff

► To cite this version:

Alessandra Paganotto, Camille Langlade, Emilie Rosset, S. Buff. Dispelling myths on external factors affecting post-thawing motility of canine spermatozoa: a retrospective study on 1242 samples. 9th International Symposium on Canine and Feline Reproduction, ICAR 2022 Satellite & 24th EVSSAR Congress, Jun 2022, Milano, Italy. hal-03720243

HAL Id: hal-03720243

<https://vetagro-sup.hal.science/hal-03720243>

Submitted on 15 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



VetAgro Sup

Dispelling myths on external factors affecting post-thawing motility of canine spermatozoa: a retrospective study on 1242 samples

A. Paganotto, C. Langlade, E. Rosset, S. Buff

Université de Lyon, VetAgro Sup, Marcy l'Etoile, France — alessandra.paganotto@vetagro-sup.fr



Introduction

Although the characteristics of canine ejaculates have been the subject of a number of studies, the literature on the **influence of donor traits and external factors on post-thawing outcome** remains scarce [1], in particular regarding the possibility of **repeating semen collection** twice in the same day. The purpose of this retrospective study was to analyse characteristics influencing post-thawing motility.

Material and methods

This analysis is based on 1,242 ejaculates frozen at VetAgro Sup between 1995 and 2021. Records included:

DONOR TRAITS AND EXTERNAL PARAMETERS

- Spermatozoa **concentration**, **motility** and **morphology** evaluated at the time of collection and 4 weeks after, **age** and dog **weight**, straw **size**, whether the collection was the **first or second** of the day and **season**.
- Depending on spermatozoa concentration, the ejaculate was divided into 0,25 or 0,5 mL straws.
- All samples were cryopreserved with a tris-egg yolk and 6% glycerol extender and stored following the same protocol [2]. Thawing control was carried out by placing the straw in a water bath (37°C) for 45 s and motility was analysed after 5 min by at least 2 different clinicians.

STATISTICAL ANALYSIS

- The influence of external parameters on motility loss was evaluated with ANOVA and Pearson's r for group differences and variable relationships respectively.

Data Analysis

The number of collections performed on the same day was only found significantly correlated with the **number of spermatozoa** ($p < 0.001$), with a lower presence for the second ejaculate [See Figure 1]. However:

- the collection repetition was not significantly correlated with post-thawing motility loss. In fact, motility loss did not differ among any of the analysed donor-related and external parameters, ie. age, weight, season, straw size
- Among the most represented races (N>30), the ones with significantly higher motility loss were the Leonberger and the Great Dane.




Figure 1:

(a) Boxplot of motility loss for most represented races (N>30).

(b) Boxplot of motility loss with respect to seasons. No significant difference was found ($p = <<<< 0,52$).

(c) Boxplot of spermatozoid number on 1st and 2nd collection. Significant mean difference was found ($p = 1,1e-15$) with a mean loss of 30% on the second collection.

(d) Boxplot of motility loss on 1st and 2nd collection. No significant difference was found ($p = 0,39$).

(e) Pearson's correlation coefficients (p -value < 0,005) between pairwise variables. *** p -value < 0,0005. ** p -value < 0,005

DISCUSSION

Recent work on ejaculates collected at short intervals for one cryopreservation process noted that their combination allows a growth in quantity without compromising semen quality [3].

We further observed in our study that the **second ejaculate** can also be utilised independently **without a significant change in post-thawing motility**. In addition, we found that neither age, time of the year, dog's weight, straw size or sperm morphology influenced post-thawing motility loss. Semen cryopreservation in dogs can therefore be performed successfully without these parameters' influence on the outcome.

References

- Lechner D, Aurich J, Schäfer-Somi S, Aurich C. Effects of age, size and season on cryotolerance of dog semen - A retrospective analysis. *Anim Reprod Sci.* 2022
- Rota, A., Linde-Forsberg, C., Vannozzi, J., Romagnoli, S., & Rodriguez-Martinez, H. Cryosurvival of Dog Spermatozoa at Different Glycerol Concentrations and Freezing/Thawing Rates. *Reproduction in Domestic Animals.* 1998
- Lechner D, Aurich J, Schäfer-Somi S, Herbel J, Aurich C. Combined cryopreservation of canine ejaculates collected at a one-hour interval increases semen doses for artificial insemination without negative effects on post-thaw sperm characteristics. *Reprod Domest Anim.* 2021



European Veterinary Society
For Small Animal Reproduction

XXIV International Congress – Milan
(IT) – June 30th to July 2nd, 2022